

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ALTUVIII0 safely and effectively. See full prescribing information for ALTUVIII0.

ALTUVIII0™ [antihemophilic factor (recombinant), Fc-VWF-XTEN fusion protein-ehtl], lyophilized powder for solution, for intravenous use

Initial U.S. Approval: 2023

INDICATIONS AND USAGE

ALTUVIII0 [antihemophilic factor (recombinant), Fc-VWF-XTEN fusion protein-ehtl] is a recombinant DNA-derived, Factor VIII concentrate indicated for use in adults and children with hemophilia A (congenital factor VIII deficiency) for:

- Routine prophylaxis to reduce the frequency of bleeding episodes
- On-demand treatment & control of bleeding episodes
- Perioperative management of bleeding (1)

Limitation of Use:

ALTUVIII0 is not indicated for the treatment of von Willebrand disease. (1)

DOSAGE AND ADMINISTRATION

For intravenous use only.

- Each ALTUVIII0 vial label states Factor VIII activity in international units (IU or unit). (2.1)
- For routine prophylaxis: 50 IU/kg once weekly. (2.1)
- For on-demand treatment and control of bleeding episodes and perioperative management: 50 IU/kg (2.1)
Estimated Increment of Factor VIII (IU/dL or % of normal) = 50 IU/kg × 2 (IU/dL per IU/kg). (2.1)

To achieve a specific target Factor VIII activity level, use the following formula: Dosage (IU) = Body Weight (kg) × Desired Factor VIII Increase (IU/dL or % normal) × 0.5 (IU/kg per IU/dL). (2.1).

DOSAGE FORMS AND STRENGTHS

For injection: nominally 250, 500, 750, 1000, 2000, 3000, or 4000 IU, lyophilized powder in single-dose vials for reconstitution. (3)

CONTRAINDICATIONS

Do not use in patients who have had severe hypersensitivity reactions, including anaphylaxis, to ALTUVIII0 or excipients of ALTUVIII0. (4)

WARNINGS AND PRECAUTIONS

- Hypersensitivity reactions, including anaphylaxis, may occur. Should symptoms occur, immediately discontinue ALTUVIII0 and initiate appropriate treatment. (5.1)
- Neutralizing antibodies (inhibitors) to Factor VIII have been reported. If expected plasma Factor VIII activity levels are not attained, or if bleeding is not controlled with an appropriate dose, perform an assay that measures Factor VIII inhibitor concentration. (5.2, 5.3)

ADVERSE REACTIONS

Most common adverse reaction (incidence >10%) are headache and arthralgia. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Bioerativ Therapeutics Inc. (A SANOFI COMPANY) at 1-800-633-1610 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

USE IN SPECIFIC POPULATIONS

Pediatric Use: No dosing adjustment is needed in this population. (8.4)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling

Revised: 02/2023

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Dose
- 2.2 Preparation and Reconstitution
- 2.3 Administration

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Hypersensitivity Reactions
- 5.2 Neutralizing Antibodies
- 5.3 Monitoring Laboratory Tests

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics
- 12.6 Immunogenicity

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

ALTUVIII0 [antihemophilic factor (recombinant), Fc-VWF-XTEN fusion protein-ehtl] is a von Willebrand Factor (VWF) independent recombinant DNA-derived, Factor VIII concentrate indicated for use in adults and children with hemophilia A (congenital factor VIII deficiency) for:

- Routine prophylaxis to reduce the frequency of bleeding episodes
- On-demand treatment and control of bleeding episodes
- Perioperative management of bleeding

Limitation of Use

ALTUVIII0 is not indicated for the treatment of von Willebrand disease.

2 DOSAGE AND ADMINISTRATION

For intravenous use after reconstitution only.

2.1 Dose

- Each ALTUVIII0 vial label states the Factor VIII potency in international units (IU). One IU corresponds to the Factor VIII activity contained in one milliliter of normal human plasma, as defined by the current World Health Organization (WHO) international standard for Factor VIII concentrate.
- Potency assignment for ALTUVIII0 is determined using an activated partial thromboplastin time (aPTT)-based one-stage clotting assay. It is recommended to use a validated one-stage clotting assay to measure ALTUVIII0 Factor VIII activity in plasma. The ALTUVIII0 Factor VIII activity level is overestimated by the chromogenic assay and a specific ellagic acid based aPTT reagent in one-stage clotting assay by approximately 2.5-fold [see *Warnings and Precautions* (5.3)].

For the dose of 50 IU/kg, the expected *in vivo* peak increase in Factor VIII level expressed as IU/dL (or % of normal) is estimated using the following formula:

Estimated Increment of Factor VIII (IU/dL or % of normal) = 50 IU/kg × 2 (IU/dL per IU/kg)

To achieve a specific target Factor VIII activity level, use the following formula: Dosage (IU) = Body Weight (kg) × Desired Factor VIII Increase (IU/dL or % normal) × 0.5 (IU/kg per IU/dL).

Routine Prophylaxis

The recommended dosing for routine prophylaxis for adults and children is 50 IU/kg of ALTUVIII0 administered once weekly.

On-demand Treatment and Control of Bleeding Episodes

ALTUVIII0 dosing for the on-demand treatment and control of bleeding episodes is provided in Table 1.

Table 1: Dosing for On-demand Treatment and Control of Bleeding Episodes

Type of Bleeding	Recommended Dose	Additional Information
Minor and Moderate For example: uncomplicated joint bleeds, minor muscular bleeds, mucosal or subcutaneous bleeds	Single dose of 50 IU/kg	For minor and moderate bleeding episodes occurring within 2 to 3 days after a prophylactic dose, a lower dose of 30 IU/kg dose may be used. Additional doses of 30 or 50 IU/kg every 2 to 3 days may be considered.
Major For example: Intracranial, retroperitoneal, iliopsoas and neck bleeds, muscle bleeds with compartment syndrome and bleeds associated with a significant decrease in the hemoglobin level	Single dose of 50 IU/kg	Additional doses of 30 or 50 IU/kg every 2 to 3 days can be considered.

For resumption of prophylaxis (if applicable) after treatment of a bleed, it is recommended to allow an interval of at least 72 hours between the last 50 IU/kg dose for treatment of a bleed and resuming prophylaxis dosing. Thereafter, prophylaxis can be continued as usual on the patient's regular schedule.

Perioperative Management

ALTUVIII0 dosing for perioperative management is provided in Table 2.

Table 2: Dosing for Perioperative Management


Type of Surgery	Pre-operative Dose	Post-operative Dose
Minor For example: Tooth extraction	Single dose of 50 IU/kg	An additional dose of 30 or 50 IU/kg after 2 to 3 days may be considered.
Major For example: Intracranial, intra-abdominal, joint replacement surgery, or complicated dental procedures.	Single dose of 50 IU/kg	Additional doses of 30 or 50 IU/kg every 2 to 3 days may be administered as clinically needed for perioperative management.

2.2 Preparation and Reconstitution


1. Use aseptic technique and a flat work surface during the reconstitution procedure.
2. Allow the ALTUVIIIIO vial, containing the white to off-white lyophilized powder, and the prefilled diluent syringe to reach room temperature before use.
3. Remove the plastic cap from the ALTUVIIIIO vial and wipe the rubber stopper of the vial with an alcohol wipe. Allow the rubber stopper to dry.
4. Completely remove the backing from the vial adapter package by peeling back the lid. **Do not** remove the vial adapter from the package or touch the inside of the package of the adapter.




5. Keep the vial on a flat surface. Hold the vial with one hand and using the other hand, place the vial adapter in its package over the vial. The spike should be placed directly above the center of the rubber stopper. Push the vial adapter straight down until the spike on the vial adapter punctures the center of the vial stopper and is fully inserted.



6. Lift the package cover away from the vial adapter and throw away the cover.



7. Only use the prefilled diluent syringe provided to reconstitute the powdered medicine. Hold the plunger rod by the circular disk. Place the tip of the plunger rod into the end of the prefilled diluent syringe. Turn the plunger rod to the right until it is firmly attached.



8. With one hand, hold the prefilled diluent syringe directly under the cap with the cap pointing up. Make sure you are holding the prefilled diluent syringe by the ridged part directly under the cap. **Do not** use if the cap has been removed or is not securely attached.
9. With your other hand, grasp the cap and bend it at a 90 degree angle until it snaps off. After the cap snaps off, you will see the glass tip of the prefilled diluent syringe. **Do not** touch the glass tip of the prefilled diluent syringe or the inside of the cap.
10. Be sure the vial is sitting on a flat surface. Insert the tip of the prefilled diluent syringe into the vial adapter opening. Turn the prefilled diluent syringe to the right until it is securely attached to the vial adapter.
11. Slowly push down on the plunger rod to inject all of the liquid (diluent) from the prefilled diluent syringe into the vial. The plunger rod may rise slightly afterward. This is normal.
12. With the prefilled diluent syringe still connected to the adapter, gently swirl the vial until the powder is completely dissolved. Check the solution through the vial to make sure the powder is fully dissolved. The solution should look clear and colorless to opalescent. **Do not** shake. **Do not** use the reconstituted ALTUVIIIIO if it contains visible particles or is cloudy.
POOLING: pooling is the process of combining two or more reconstituted vials into a larger luer lock syringe (not provided in the carton). If the dose requires more than one vial, reconstitute each vial as described above (See Steps 3–12) with the prefilled diluent syringe provided. **Do not** detach the prefilled diluent syringe until you are ready to attach the large luer lock syringe to the next vial. Keep the vial adapter attached to the vial as you will need it for attaching a large luer lock syringe. Use a larger luer lock plastic syringe to combine the contents of the reconstituted vials into the syringe, similar as described in the Steps 13–14. Repeat this pooling procedure with each vial you will be using. Once you have pooled the required dose, proceed with the administration steps using the larger luer lock syringe.
13. Make sure the plunger rod is pressed all the way down and the diluent syringe is firmly attached to the vial adapter. Turn the vial upside-down. Slowly pull down on the plunger rod to draw all the solution from the vial into the diluent syringe. Be careful not to pull the plunger rod completely out of the diluent syringe.
14. Gently unscrew the diluent syringe from the vial adapter by turning it to the right. Dispose of the vial with the adapter still attached. If you are not ready to inject, put the syringe cap carefully back onto the syringe tip. **Do not** touch the syringe tip or the inside of the cap.
15. Use the reconstituted ALTUVIIIIO as soon as possible, but no later than **3 hours** after reconstitution. Do not touch the glass tip of the syringe if not used immediately after reconstitution. Protect from direct sunlight. Do not refrigerate after reconstitution.

2.3 Administration

For intravenous use only.

- Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. **Do not** use ALTUVIIIIO solution if particulate matter or discoloration is observed.
- **Do not** administer reconstituted ALTUVIIIIO in the same tubing or container with other medications.

Administration Steps:

1. Attach the syringe to the connector end of the infusion set tubing by turning it to the right until it is securely attached.
2. Push the plunger rod until all air is removed from the syringe and ALTUVIIIIO has filled the infusion set needle. **Do not** push ALTUVIIIIO solution through the needle.
3. Remove the protective needle cover from the infusion set needle.
4. Perform intravenous injection. The rate of administration should be determined by the patient's comfort level and no faster than:
 - For adults and adolescents: 1–2 minutes per vial.
 - For children: 2–3 minutes per vial if body weight is greater than or equal to 20 kg; or 6 minutes per vial if body weight is less than 20 kg.
5. After infusing ALTUVIIIIO, remove and properly discard the infusion set.

3 DOSAGE FORMS AND STRENGTHS

ALTUVIIIIO is available as a white to off-white lyophilized powder for reconstitution in single-dose vials containing nominally 250, 500, 750, 1000, 2000, 3000, or 4000 international units (IU) per vial.

4 CONTRAINDICATIONS

ALTUVIIIIO is contraindicated in patients who have had severe hypersensitivity reactions, including anaphylaxis, to the product or its excipients [see Description (11)].

5 WARNINGS AND PRECAUTIONS

5.1 Hypersensitivity Reactions

Allergic-type hypersensitivity reactions, including anaphylaxis, may occur with ALTUVIIIIO. Allergic-type hypersensitivity reactions were not reported in the clinical trials. Inform patients of signs of hypersensitivity reactions that may progress to anaphylaxis (including hives, shortness of breath, chest tightness, wheezing, hypotension and itching). Advise patients to discontinue use of ALTUVIIIIO if hypersensitivity symptoms occur and contact a physician and/or seek immediate emergency care.

5.2 Neutralizing Antibodies

Formation of neutralizing antibodies (inhibitors) to Factor VIII are possible following administration of ALTUVIIIIO. Neutralizing antibodies were not reported in the clinical trials. Monitor all patients for the development of Factor VIII inhibitors by appropriate clinical observations and laboratory tests. If the patient's plasma Factor VIII level fails to increase as expected or if bleeding is not controlled after ALTUVIIIIO administration, the presence of an inhibitor (neutralizing antibodies) should be suspected, and appropriate testing performed [see Warnings and Precautions (5.3)].

5.3 Monitoring Laboratory Tests

If assessment of plasma Factor VIII activity is needed, it is recommended to use a validated one-stage clotting assay [see Dosage and Administration (2)]. The ALTUVIIIIO Factor VIII activity level is overestimated by the chromogenic assay and a specific ellagic acid based aPTT reagent in one-stage clotting assay by approximately 2.5-fold. If these assays are used, divide the result by 2.5 to approximate the patient's ALTUVIIIIO Factor VIII activity level. Use of a reference laboratory is recommended when a qualified one-stage clotting assay or chromogenic assay is not available locally. Monitor for the development of Factor VIII inhibitors. If bleeding is not controlled with ALTUVIIIIO and the expected factor VIII activity plasma levels are not attained, perform an assay to determine if Factor VIII inhibitors are present (use Bethesda Units to titer inhibitors).

6 ADVERSE REACTIONS

The most common adverse reactions (>10% of subjects) reported in clinical trials were headache and arthralgia.

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The safety of ALTUVIII0 has been evaluated in 159 previously treated patients (PTPs) (134 adults and 25 adolescents) with severe Hemophilia A (<1% endogenous FVIII activity or a genetic mutation consistent with severe Hemophilia A) who received at least one dose of ALTUVIII0 for either routine prophylaxis, on-demand treatment of bleeding episodes or perioperative management. A total of 152 (96%) subjects achieved at least 25 exposure days and 115 (72%) subjects achieved at least 50 exposure days with a median of 53.0 (range 2–63) for both exposure days and injections per subject. Overall exposure was monitored for a total of 151.5 subject-years. Adverse drug reactions (ADRs) (summarized in Table 3) were reported in 57 (36%) of the 159 subjects treated with routine prophylaxis or on-demand therapy. There were no age-specific differences in ADRs observed between adolescent and adult subjects. In the study, no inhibitors were detected and no ADRs of anaphylaxis were reported.

Table 3: Adverse Drug Reactions with Frequency of ≥3% Reported for ALTUVIII0

MedDRA System Organ Class	Adverse Drug Reactions	Number of Subjects n (%) (N = 159)
Nervous system disorders	Headache*	33 (21)
Musculoskeletal and connective tissue disorders		
	Arthralgia	26 (16)
	Back pain	9 (6)

*Includes preferred terms of headache (32 subjects) and migraine (1 subject).

In the ongoing pediatric study, the safety of ALTUVIII0 was evaluated in 67 male PTPs <12 years of age with severe hemophilia A who received at least one dose of ALTUVIII0. At the time of the interim analysis, a total of 23 (34%) subjects achieved at least 25 exposure days with a median of 14.0 (range 1–46) for both exposure days and injections per subject.

Adverse drug reactions, headache, were reported in 1 (1%) of subjects. In the study, no inhibitors were detected and no ADRs of anaphylaxis were reported.

Thromboembolic events occurred in 1% (3/206) of subjects in the long term safety extension study; these three subjects had pre-existing risk factors.

Immunogenicity

All subjects were monitored for neutralizing antibodies (inhibitors) to Factor VIII in the clinical program. No subjects developed neutralizing antibodies to Factor VIII [see Immunogenicity (12.6)].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

There are no data with ALTUVIII0 use in pregnant women to inform a drug-associated risk. Animal developmental and reproductive studies have not been conducted with ALTUVIII0. Therefore, it is not known whether ALTUVIII0 can affect reproductive capacity or cause fetal harm when given to pregnant women.

In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2–4% and 15–20%, respectively.

8.2 Lactation

Risk Summary

There is no information regarding the presence of ALTUVIII0 in human milk, its effects on the breastfed infant, or its effects on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for ALTUVIII0 and any potential adverse effects on the breastfed infant from ALTUVIII0 or from the underlying maternal condition.

8.4 Pediatric Use

Safety, efficacy, and pharmacokinetics (PK) have been evaluated in 92 previously treated, pediatric patients <18 years of age who received at least one dose of ALTUVIII0 as part of routine prophylaxis, treatment of bleeding episodes, or perioperative management. Adolescent subjects were enrolled in the adult and adolescent study and pediatric subjects <12 years of age were enrolled in an ongoing pediatric trial. Thirty-one subjects (33.7%) were <6 years of age, 36 (39.1%) subjects were 6 to <12 years of age, and 25 subjects (27.2%) were adolescents (12 to <18 years of age). Interim data from a pediatric study of the 67 subjects <12 years of age showed that no dosing adjustment was required [see Clinical Pharmacology (12)].

8.5 Geriatric Use

Clinical studies of ALTUVIII0 did not include sufficient numbers of subjects 65 years of age and older to determine whether or not they respond differently from younger subjects. However, clinical experience with other Factor VIII products has not identified differences between the elderly and younger patients.

11 DESCRIPTION

ALTUVIII0 [antihemophilic factor (recombinant), Fc-VWF-XTEN fusion protein-ehtl] is a sterile, non-pyrogenic, white to off-white lyophilized powder for reconstitution for intravenous injection. The product is supplied in single-dose vials containing nominal potencies of 250, 500, 750, 1000, 2000, 3000, or 4000 international units (IU). Each vial of ALTUVIII0 is labeled with the actual Factor VIII activity content in IU. The powder for injection is reconstituted with 3 mL sterile water for injection (SWFI) supplied in a sterile pre-filled syringe. The reconstituted solution should be essentially free of particles. The final product contains the excipients: arginine hydrochloride (250 mM), calcium chloride dihydrate (5 mM), histidine (10 mM), polysorbate 80 (0.05% w/v), and sucrose (5% w/v).

The active ingredient in ALTUVIII0 is a fully recombinant fusion protein comprising a single chain B-domain deleted (BDD) analogue of human FVIII covalently fused to the Fc domain of human immunoglobulin G1 (IgG1), the FVIII-binding D'D3 domain of human von Willebrand factor (VWF), and 2 XTEN polypeptides. ALTUVIII0 contains 2829 amino acids with an apparent molecular weight of 312 kDa. ALTUVIII0 is synthesized as 2 polypeptide chains which are covalently linked by 2 Fc hinge disulfide bonds. The first FVIII-XTEN-Fc polypeptide chain contains the A1A2 domain of FVIII along with 5 amino acids from B-domain (1–745 amino acids) fused to the 288-XTEN polypeptide (in place of the natural FVIII B-domain), the A3C1C2 domain of FVIII (1649–2332), and the Fc domain of human IgG1. The second VWF-XTEN-a2-Fc polypeptide chain contains the D'D3 domain of VWF (1–477 amino acids) fused to the 144-XTEN polypeptide, a thrombin cleavable acidic region 2 sequence from FVIII and the Fc domain of human IgG1. The Fc domain includes the hinge, CH₂, and CH₃ domains of IgG1. The Fc, VWF, and XTEN portions of the molecule extend the half-life of ALTUVIII0 in plasma. ALTUVIII0 is produced by recombinant DNA technology in a human embryonic kidney (HEK) cell line, which has been extensively characterized. ALTUVIII0 is manufactured without addition of human- or

animal-derived components and purified by a combination of multiple chromatography steps, a detergent viral inactivation step, a nano filtration step for viral clearance, and ultrafiltration steps.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

ALTUVIII0 [antihemophilic factor (recombinant), Fc-VWF-XTEN fusion protein-ehtl] temporarily replaces the missing coagulation factor VIII needed for effective hemostasis. ALTUVIII0 has demonstrated 3- to 4-fold prolonged half-life relative to other standard and extended half-life FVIII products.

Mechanism of Half-life Extension

ALTUVIII0 is a recombinant FVIII analogue fusion protein that is independent of endogenous VWF in order to overcome the half-life limit imposed by FVIII-VWF interactions. The D'D3 domain of VWF is the region that interacts with FVIII. Appending the D'D3 domain of VWF to a recombinant FVIII-Fc fusion protein provides protection and stability to FVIII, and prevents FVIII interaction with endogenous VWF, thus overcoming the limitation on FVIII half-life imposed by VWF clearance.

The Fc region of human immunoglobulin G1 (IgG1) binds to the neonatal Fc receptor (FcRn). FcRn is part of a naturally occurring pathway that delays lysosomal degradation of immunoglobulins by recycling them back into circulation, thus prolonging the plasma half-life of the fusion protein.

ALTUVIII0 contains 2 XTEN polypeptides, which alter the hydrodynamic radius of the fusion protein, thus reducing rates of clearance and degradation, and improving pharmacokinetic properties. In ALTUVIII0, the natural FVIII B domain (except 5 amino acids) is replaced with the first XTEN, inserted in between FVIII N745 and E1649 amino acid residues; and the second XTEN is inserted in between the D'D3 domain and Fc.

12.2 Pharmacodynamics

Hemophilia A is a bleeding disorder characterized by a deficiency of functional coagulation factor VIII (FVIII), which leads to a prolonged clotting time in the activated partial thromboplastin time (aPTT)-based one-stage clotting assay. Administration of ALTUVIII0 increases plasma levels of FVIII, temporarily correcting the coagulation defect in hemophilia A patients.

Based on FVIII pharmacokinetic/pharmacodynamic analyses, the risk of bleeding is negatively correlated with FVIII activity. Once weekly 50 IU/kg ALTUVIII0 provided factor VIII activity levels that were associated with a low bleed risk.

12.3 Pharmacokinetics

The PK of ALTUVIII0 were evaluated in prospective, open-label clinical studies, enrolling 159 adults and adolescents, and 67 children <12 years old, respectively, receiving weekly IV injection of 50 IU/kg. Among children <12 years old, 32 subjects had ALTUVIII0 single dose PK profiles available.

PK parameters following a single dose of ALTUVIII0 are presented in Table 4. The PK parameters were based on plasma FVIII activity measured by the aPTT-based one-stage clotting assay. After a single dose of 50 IU/kg, ALTUVIII0 exhibited high sustained FVIII activity with prolonged half-life across age cohorts. There was a trend of increasing area under the curve (AUC), and decreasing clearance, with increasing age in the pediatric cohorts. The PK profile at steady state (Week 26) was comparable with the PK profile obtained after the first dose.

Table 4: Pharmacokinetic Parameters Following a Single Dose of ALTUVIII0 by age (one-stage clotting assay)

PK Parameters (mean SD)	Pediatric Study	Pediatric Study	Adult and Adolescent Study	Adult and Adolescent Study
	1 to <6 Years N = 14	6 to <12 Years N = 18	12 to <18 years N = 25	Adults N = 134
AUC (IU·h/dL)	6710 (1190)	7190 (1450)	8350 (1550)	9850 (2010) *
t _{1/2} (h)	39.9 (5.71)	42.4 (3.70)	44.6 (4.99)	48.2 (9.31)
CL (mL/h/kg)	0.740 (0.128)	0.681 (0.139)	0.582 (0.115)	0.493 (0.121) *
V _{ss} (mL/kg)	38.0 (7.19)	38.1 (6.80)	34.9 (7.38)	31.0 (7.32) *
MRT (hr)	51.9 (9.06)	56.3 (5.10)	60.0 (5.54)	63.9 (10.2) *

AUC_{0–tau} = area under the activity-time curve over the dosing interval, CL = clearance, MRT = mean residence time, SD = standard deviation, t_{1/2z} = terminal half-life, V_{ss} = volume of distribution at steady state.

*Calculation based on 128 profiles.

ALTUVIII0 at steady state maintained normal to near normal (>40 IU/dL) FVIII activity for a mean (SD) of 4.1 (0.7) days with once weekly prophylaxis in adults. The FVIII activity over 10 IU/dL was maintained in 83.5% of adults and adolescent subjects throughout the study. In children <12 years ALTUVIII0 maintained normal to near normal (>40 IU/dL) FVIII activity for 2 to 3 days and >10 IU/dL FVIII activity for approximately 6 to 7 days. The majority of children <12 years maintained FVIII activity in mild hemophilia range (>5 IU/dL) 7 days after the dosing (see Table 5).

Table 5: Pharmacokinetic Parameters at Steady State of ALTUVIII0 by age (one-stage clotting assay)

PK Parameters Mean (SD)	Pediatric Study	Pediatric Study	Adult and Adolescent Study	Adult and Adolescent Study
	1 to <6 years N = 20	6 to <12 years N = 35	12 to <18 years N = 24	Adults N = 124
Peak (IU/dL)	113 (26.2)	121 (25.31)	124 (31.2)	150 (35.0)
IR (kg×IU/dL/IU)	2.10 (0.53)	2.17 (0.46)	2.25 (0.61)	2.64 (0.61)
Time to 40 IU/dL (h) †	59.2 (7.68)	72.2 (11.2)	81.7 (13.1)	97.0 (20.2)
Time to 20 IU/dL (h) †	99.3 (9.15)	117 (13.9)	130 (15.7)	150 (27.7)

Table 5: Pharmacokinetic Parameters at Steady State of ALTUVIII0 by age (one-stage clotting assay) (continued)

PK Parameters Mean (SD)	Pediatric Study	Pediatric Study	Adult and Adolescent Study [*]	Adult and Adolescent Study [*]
	1 to <6 years N = 20	6 to <12 years N = 35	12 to <18 years N = 24	Adults N = 124
Time to 10 IU/dL (h) †	139 (11.7)	163 (17.1)	179 (21.4)	200 (35.4)
Trough (IU/dL)	6.75 (2.29)	9.77 (3.64)	9.23 (4.77)	18.0 (16.6)

Peak = 15 min post dose at steady state, IR = incremental recovery, Trough – predose FVIII activity value at steady state, SD = standard deviation.

*Steady state peak, trough and IR were computed using available measurements at week 4 and week 52/End of study PK sampling visit for pediatric and adult and adolescent studies, respectively.

† Time to FVIII activity was predicted using population PK model.

Specific Populations

The following factors have no clinically meaningful effect on the pharmacokinetics of ALTUVIII0: age (1.4 to 72 years), sex, race (White, Asian), VWF antigen activity (40 to 339 IU/dL), hematocrit level (28% to 57%), blood type, HCV status, or HIV status. Body weight (12.5 to 133 kg) is expected to alter weight normalized clearance (dL/h/kg) by 79% to -18% compared to a typical patient.

12.6 Immunogenicity

The observed incidence of anti-drug antibodies (ADAs) is highly dependent on the sensitivity and specificity of the assay. Differences in assay methods preclude meaningful comparisons of the incidence of anti-drug antibodies in the studies described below with the incidence of anti-drug antibodies in other studies, or of other Factor VIII products.

All subjects were monitored for neutralizing antibodies (inhibitors) to Factor VIII in the clinical program. No subjects developed neutralizing antibodies to Factor VIII.

During treatment (up to 49.64 weeks) in the clinical studies, 4/277 (2.2%) of ALTUVIII0 patients developed anti-drug antibodies.

No impact of ADAs on the FVIII activity-time profiles and PK exposure parameters was observed. No impact of ADAs with respect to bleeding episodes and in the pharmacodynamic response was noted.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

No carcinogenicity or mutagenicity studies have been conducted with ALTUVIII0. No studies have been conducted to evaluate the effects of ALTUVIII0 on fertility.

14 CLINICAL STUDIES

The safety, efficacy, and pharmacokinetics of ALTUVIII0 were evaluated in two multicenter, prospective, open-label clinical studies (one study in adults and adolescents ≥12 years of age and one pediatric study in children <12 years of age) in previously treated patients (PTPs) with severe hemophilia A (<1% endogenous Factor VIII activity or a documented genetic mutation consistent with severe hemophilia A).

All studies evaluated the efficacy of routine prophylaxis with a weekly dose of 50 IU/kg and determined hemostatic efficacy in the treatment of bleeding episodes and during perioperative management in subjects undergoing major or minor surgical procedures.

The completed adult and adolescent study enrolled a total of 159 PTPs (158 male and 1 female subjects) with severe hemophilia A. Subjects were aged 12 to 72 years and included 25 adolescent subjects aged 12 to 17 years. All 159 enrolled subjects received at least one dose of ALTUVIII0 and were evaluable for efficacy. A total of 149 subjects (93.7%) completed the study.

The ongoing pediatric study enrolled 67 male PTPs <12 years of age with severe hemophilia A (31 subjects were 1 to 5 years of age and 36 were 6 to 11 years of age) at data cut-off. Of the 67 enrolled subjects, all received at least 1 dose of ALTUVIII0.

Routine Prophylaxis to Reduce Bleeding Episodes

Adult and Adolescent Study

The efficacy of weekly 50 IU/kg ALTUVIII0 as routine prophylaxis was evaluated as estimated by the mean annualized bleed rate (ABR) and by comparing the ABR during on-study prophylaxis vs. the ABR during pre-study FVIII prophylaxis. A total of 133 adults and adolescents, who were on pre-study FVIII prophylaxis, were assigned to receive ALTUVIII0 for routine prophylaxis at a dose of 50 IU/kg IV once weekly for 52 weeks (Arm A). An additional 26 subjects, who were on pre-study episodic (on-demand) treatment with FVIII, received episodic (on-demand) treatment with ALTUVIII0 at doses of 50 IU/kg IV for 26 weeks, followed by routine prophylaxis at a dose of 50 IU/kg IV once weekly for 26 weeks (Arm B). Overall, 115 subjects received at least a total number of 50 exposure days (EDs) in Arm A and 17 subjects completed at least 25 EDs of routine prophylaxis in Arm B.

The ABR in subjects evaluable for efficacy with at least 26 weeks of exposure are summarized in Table 6. Routine prophylaxis resulted in a mean ABR (95% CI) of 0.7 (0.5, 1.0), a median (Q1, Q3) ABR of 0 (0, 1.0), and a median (Q1, Q3) annualized joint bleeding rate of 0 (0, 1.0).

Table 6: Summary of Annualized Bleeding Rate (ABR) with ALTUVIII0 Prophylaxis, ALTUVIII0 On-demand Treatment, and After Switch to ALTUVIII0 Prophylaxis in Patients ≥12 Years of Age

Endpoint [*]	Arm A Prophylaxis [†]	Arm B On-demand [‡]	Arm B Prophylaxis [‡]
	N = 128	N = 26	N = 26
Treated bleeds			
Mean ABR (95% CI) [§]	0.7 (0.5, 1.0)	21.4 (18.8, 24.4)	0.7 (0.3, 1.5)
Median ABR (Q1, Q3)	0 (0, 1.0)	21.1 (15.1, 27.1)	0 (0, 0)
subjects with zero bleeds, n (%)	82 (64.1)	0	20 (76.9)

Table 6: Summary of Annualized Bleeding Rate (ABR) with ALTUVIII0 Prophylaxis, ALTUVIII0 On-demand Treatment, and After Switch to ALTUVIII0 Prophylaxis in Patients ≥12 Years of Age (continued)

Endpoint [*]	Arm A Prophylaxis [†]	Arm B On-demand [‡]	Arm B Prophylaxis [‡]
Treated spontaneous bleeds			
Mean ABR (95% CI) [§]	0.3 (0.2, 0.4)	15.8 (12.3, 20.4)	0.4 (0.2, 1.2)
Median ABR (Q1, Q3)	0 (0, 0)	16.7 (8.6, 23.8)	0 (0, 0)
% subjects with zero bleeds, n (%)	103 (80.5)	1 (3.8)	22 (84.6)
Treated joint bleeds			
Mean ABR (95% CI) [§]	0.5 (0.4, 0.7)	17.5 (14.9, 20.5)	0.6 (0.3, 1.5)
Median ABR (Q1, Q3)	0 (0, 1.0)	18.4 (10.8, 23.9)	0 (0, 0)
% subjects with zero bleeds, n (%)	92 (71.9)	0	21 (80.8)
All Bleeds (treated and untreated)[*]			
Mean ABR (95% CI) [§]	1.1 (0.8, 1.5)	22.2 (19.4, 25.4)	0.9 (0.4, 1.8)
Median ABR (Q1, Q3)	0 (0, 1.2)	21.1 (16.8, 27.1)	0 (0, 1.9)
% subjects with zero bleeds, n (%)	71 (55.5)	0	19 (73.1)

ABR = annualized bleed rate; CI = confidence interval; Q1= 25th percentile, Q3=75th percentile.

*Reflects all bleeds reported by patients including those where no ALTUVIII0 was administered.

†Subjects assigned to receive ALTUVIII0 prophylaxis for 52 weeks.

‡Subjects assigned to receive ALTUVIII0 for 26 weeks.

§Based on negative binomial model.

An intra-subject comparison (N = 78) between mean ABR during on-study prophylaxis with ALTUVIII0 and that during pre-study FVIII prophylaxis yielded a 77% reduction in treated bleeds (95% CI: 58%, 87%).

All subjects with target joints at baseline (defined as ≥3 spontaneous bleeding episodes in a major joint which occurred in a consecutive 6-month period) achieved resolution of all target joints (45/45, 100%) with 12 months of prophylactic treatment with ALTUVIII0 (defined as ≤2 bleeding episodes in the target joint in 12 months).

Pediatric Study

The efficacy of weekly 50 IU/kg ALTUVIII0 as routine prophylaxis in children <12 years was evaluated as estimated by the mean annualized bleed rate (ABR). At the time of the interim analysis, a total of 67 children (31 children <6 years of age and 36 children 6 to <12 years of age) were enrolled to receive ALTUVIII0 for routine prophylaxis at a dose of 50 IU/kg IV once weekly for 52 weeks. In subjects with at least 26 weeks of exposure (N=23), routine prophylaxis resulted in a mean ABR (95% CI) of 0.5 (0.2, 1.3) and a median (Q1, Q3) ABR of 0 (0, 1.3) for treated bleeds. For all bleeds (treated and non treated), the mean ABR (95% CI) was 3.6 (1.6, 8.4) and the median (Q1, Q3) ABR was 0 (0, 4.5).

Efficacy in Control of Bleeding

In the adult and adolescent study, a total of 362 bleeding episodes were treated with ALTUVIII0, most occurring during on-demand treatment in Arm B. Majority of bleeding episodes were localized in joints. Response to the first injection was assessed by subjects at least 8 hours after treatment. A 4-point rating scale of excellent, good, moderate, and no response was used to assess response. Bleeding was resolved with a single 50 IU/kg injection of ALTUVIII0 in 96.7% of bleeding episodes. The median (Q1; Q3) total dose to treat a bleeding episode was 50.9 IU/kg (50.0; 51.9). Control of bleeding episodes was similar across the treatment arms.

Perioperative Management of Bleeding

Perioperative hemostasis was assessed in 13 major surgeries in 12 subjects (11 adults and 1 child). Of the 13 major surgeries, 12 surgeries required a single pre-operative dose to maintain hemostasis during surgery; for 1 major surgery during routine prophylaxis no pre-operative loading dose was administered on the day of/or before surgery. The median dose per pre-operative injection was 49.96 IU/kg (range 12.7 – 61.9).

The clinical evaluation of hemostatic response during major surgery was assessed using a 4-point scale of excellent, good, moderate, or poor/none. The hemostatic effect of ALTUVIII0 was rated as “excellent” in 13 of 13 surgeries (100%). No surgery had an outcome rated as “poor/none” or “missing.”

Types of major surgeries assessed include major orthopedic procedures such as joint arthroplasties (joint replacements of knee, hip, and elbow), joint revisions and ankle fusion. Other major surgeries included molar extractions and rhinoplasty/mentoplasty.

Perioperative hemostasis was assessed in 22 minor surgeries in 19 subjects (12 adults and 7 children). The hemostatic response was evaluated by the investigator/surgeon in 15 of these minor surgeries; an excellent response was reported in all (100%).

16 HOW SUPPLIED/STORAGE AND HANDLING

How Supplied

ALTUVIII0 is supplied in kits comprising a single-dose vial containing nominally, 250, 500, 750, 1000, 2000, 3000, or 4000 international units (IU) of Factor VIII potency, a prefilled syringe with 3 mL sterile water for injection, and a sterile vial adapter (reconstitution device). The actual amount of ALTUVIII0 in IU is stated on the label and carton of each vial.

Not made with natural rubber latex.

Strength	Potency Color Code	Kit NDC Number
250 IU	Yellow	71104-978-01
500 IU	Red	71104-979-01

(continued)

Strength	Potency Color Code	Kit NDC Number
750 IU	Garnet	71104-980-01
1000 IU	Green	71104-981-01
2000 IU	Royal Blue	71104-982-01
3000 IU	Mist Grey	71104-983-01
4000 IU	Orange	71104-984-01

Not all pack sizes may be marketed.

Storage and Handling

Prior to reconstitution:

- Store ALTUVIIIIO in the original package to protect the ALTUVIIIIO vials from light.
- Store ALTUVIIIIO in powder form at 2°C to 8°C (36°F to 46°F). Do not freeze to avoid damage to the pre-filled diluent syringe.
- ALTUVIIIIO may be stored at room temperature, not to exceed 30°C (86°F), for a single period of up to 6 months, within the expiration date printed on the label.
- If stored at room temperature, record the date that ALTUVIIIIO is removed from refrigeration on the carton in the area provided. After storage at room temperature, do not return the product to the refrigerator.
- Do not use beyond the expiration date printed on the vial or 6 months after the date that was written on the carton, whichever is earlier.

After Reconstitution:

- The reconstituted product may be stored at room temperature, not to exceed 30°C (86°F), for up to 3 hours. Protect from direct sunlight. After reconstitution, if the product is not used within 3 hours, it must be discarded.
- Do not use ALTUVIIIIO if the reconstituted solution is cloudy or has particulate matter.
- Discard any unused ALTUVIIIIO.

17 PATIENT COUNSELING INFORMATION

Advise the patients to:

- Read the FDA-approved patient labeling (Patient Information and Instructions for Use).
- Call their healthcare provider or go to the emergency department right away if a hypersensitivity reaction occurs. Early signs of hypersensitivity reactions may include rash, hives, itching, facial swelling, tightness of the chest, and wheezing.
- Contact their healthcare provider or treatment facility for further treatment and/or assessment if they experience a lack of a clinical response to Factor VIII therapy because this may be a sign of inhibitor development.

Manufactured by:
Bioerativ Therapeutics Inc.
Waltham, MA 02451
A SANOFI COMPANY
US License Number 2078

©2023 Bioerativ Therapeutics Inc. All rights reserved.

For patent information: <https://www.sanofi.us/en/products-and-resources/patents>

ALTUVIIIIO™ is a trademark of Bioerativ Therapeutics Inc.

Patient Information

ALTUVIIIIO™ (al too'vee oh)

[antihemophilic factor (recombinant), Fc-VWF-XTEN fusion protein-eh1]

for intravenous use after reconstitution only

Single-dose vial

Please read this Patient Information carefully before using ALTUVIIIIO and each time you get a refill, as there may be new information. This Patient Information does not take the place of talking with your healthcare provider about your medical condition or your treatment.

What is the most important information I need to know about ALTUVIIIIO?

Do not attempt to give yourself an injection unless you have been taught how by your healthcare provider or hemophilia center.

You must carefully follow your healthcare provider's instructions regarding the dose and schedule for injecting ALTUVIIIIO so that your treatment will work best for you.

What is ALTUVIIIIO?

ALTUVIIIIO is an injectable medicine that is used to control and reduce the number of bleeding episodes in people with Hemophilia A (congenital Factor VIII deficiency).

Your healthcare provider may give you ALTUVIIIIO when you have surgery.

Who should not use ALTUVIIIIO?

You should not use ALTUVIIIIO if you had an allergic reaction to it in the past.

What should I tell my healthcare provider before using ALTUVIIIIO?

Talk to your healthcare provider about:

- Any medical problems that you have or had.
- All prescription and non-prescription medicines that you take, including over-the-counter medicines, supplements or herbal medicines.
- Pregnancy or if you are planning to become pregnant. It is not known if ALTUVIIIIO may harm your unborn baby.
- Breastfeeding. It is not known if ALTUVIIIIO passes into the milk and if it can harm your baby.

How should I use ALTUVIIIIO?

You get ALTUVIIIIO as an injection into your vein. Your healthcare provider will instruct you on how to do injections on your own, and may watch you give yourself the first dose of ALTUVIIIIO.

Contact your healthcare provider right away if bleeding is not controlled after using ALTUVIIIIO.

What are the possible side effects of ALTUVIIIIO?

You can have an allergic reaction to ALTUVIIIIO. Call your healthcare provider or emergency department right away if you have any of the following symptoms: difficulty breathing, chest tightness, swelling of the face, rash or hives.

Your body can also make antibodies called "inhibitors" against ALTUVIIIIO. This can stop ALTUVIIIIO from working properly. Your healthcare provider may give you blood tests to check for inhibitors.

The common side effects of ALTUVIIIIO are headache, joint pain, and back pain.

These are not the only possible side effects of ALTUVIIIIO. Tell your healthcare provider about any side effect that bothers you or does not go away.

What are the ALTUVIIIIO dosage strengths?

ALTUVIIIIO comes in seven different dosage strengths with 3 mL sterile water for injection (SWFI). The actual number of international units (IU) of Factor VIII activity in the vial will be imprinted on the label and on the box. The seven different strengths are as follows:

Strength	Cap Color
250 IU	Yellow
500 IU	Red
750 IU	Garnet
1000 IU	Green
2000 IU	Royal Blue
3000 IU	Mist Grey
4000 IU	Orange

Always check the actual dosage strength printed on the label to make sure you are using the strength prescribed by your healthcare provider.

How should I store ALTUVIIIIO?

- Keep ALTUVIIIIO in its original package.
- Protect it from light.
- Do not freeze.
- Store refrigerated 2°C to 8°C (36°F to 46°F) up to 48 months or at room temperature [not to exceed 30°C (86°F)], for a single period up to 6 months. Do not use ALTUVIIIIO after the expiration date printed on the label and carton of each vial.
- When storing at room temperature:
 - Note on the carton the date on which the product is removed from refrigeration.
 - Use the product before the end of this 6-month period or discard it.
 - Do not return the product to the refrigerator.

After mixing with the diluent:

- Do not use ALTUVIIIIO if the mixed solution is not clear and colorless to slightly yellowish.
- Use mixed product as soon as possible.
- You may store mixed ALTUVIIIIO at room temperature, not to exceed 30°C (86°F), for up to **3 hours**. Protect the mixed ALTUVIIIIO from direct sunlight. Discard any mixed ALTUVIIIIO not used within 3 hours.

What else should I know about ALTUVIIIIO?

Medicines are sometimes prescribed for purposes other than those listed here. Do not use ALTUVIIIIO for a condition for which it was not prescribed. Do not share ALTUVIIIIO with other people, even if they have the same symptoms that you have.

Manufactured by:
Bioverativ Therapeutics Inc.
Waltham, MA 02451
A SANOFI COMPANY
US License Number 2078

©2023 Bioverativ Therapeutics Inc. All rights reserved.

ALTUVIIIIO™ is a trademark of Bioverativ Therapeutics Inc.

INSTRUCTIONS FOR USE
ALTUVIIIIO™ (al too'vee oh)
[antihemophilic factor (recombinant), Fc-VWF-XTEN fusion protein-ehtl]
for intravenous use after reconstitution only
Single-dose vial

Read the Instructions for Use before you start using ALTUVIIIIO and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or your treatment.
Your healthcare provider should show you or your caregiver how to mix (reconstitute) and give ALTUVIIIIO the first time ALTUVIIIIO is used.
Any further questions? Ask your healthcare provider or call 1-800-633-1610.

Important Information You Need to Know Before Injecting ALTUVIIIIO
It is important that you do not try to inject ALTUVIIIIO unless you have received training from a healthcare provider.

- Read all of the instructions carefully before using the ALTUVIIIIO.
- ALTUVIIIIO is supplied as a single-dose vial of powder for mixing (reconstitution). Before you inject ALTUVIIIIO, you must mix (reconstitute) ALTUVIIIIO powder with liquid (diluent) that comes in a prefilled diluent syringe.
- After the powder and diluent are mixed the medicine is given in the vein (intravenous injection).
- The vial and the diluent syringe are not made with natural rubber latex.
- **Do not** use ALTUVIIIIO if it has been dropped on a hard surface or damaged.
- **Do not** use if the syringe cap has been removed or is not securely attached.
- **Do not** use the mixed ALTUVIIIIO if it contains visible particles or is cloudy.
- **Do not** give mixed ALTUVIIIIO in the same tubing or container with other medicines.

Storing ALTUVIIIIO

- Keep unused ALTUVIIIIO kit in the original carton and store in the refrigerator between 2°C and 8°C (36°F and 46°F). The product may be stored at room temperature up to 30°C (86°F). If stored at room temperature, the product (prior to mixing) expires after 6 months or after the expiration date on the product vial, whichever is earlier.
- Remove the product kit from the refrigerator and allow the ALTUVIIIIO vial and the prefilled diluent syringe to come to room temperature prior to injection.
- **Do not** use external heat sources such as putting the vial or prefilled diluent syringe in hot water.
- **Do not** return room temperature ALTUVIIIIO to refrigerator.
- Keep away from direct sunlight.
- **Do not** freeze.

Storing of Mixed (Reconstituted ALTUVIIIIO)

- ALTUVIIIIO should be given within **3 hours** after mixing.
- Keep away from direct sunlight.
- Do not refrigerate after mixing.
- Keep ALTUVIIIIO and all medicines out of the reach of children.

Preparing to Inject ALTUVIIIIO
Mixing (Reconstitution)

Step 1:



Look at the ALTUVIIIIO kit:

- Check that you have the correct medicine and dose.
- Check the expiration date.
- ▲ **Do not** use ALTUVIIIIO if the expiration date has passed.


Step 2:



Wash your hands with soap and water. Find a clean, flat work surface. Remove the supplies from the carton:

- Vial adapter in its package
 - Vial with powdered medicine
 - Plunger rod
 - Prefilled diluent syringe
- Also ensure you have the following supplies (not included in the carton):
- Infusion set
 - Tourniquet
 - 2 alcohol wipes
 - 1 cotton ball or gauze pad
 - 1 adhesive bandage, if required (See Step 22)
 - 1 tape, if required (See Step 20)
 - 1 larger luer lock syringe, if required (See Step 13)
 - FDA-cleared sharps disposal container (See Step 23)
- ▲ **Do not** use ALTUVIIIIO (vial or prefilled diluent syringe) if it has been dropped on a hard surface or damaged.


Step 3:



Allow the ALTUVIIIIO vial and the prefilled diluent syringe to come to room temperature before use.

- ▲ **Do not** use external heat sources such as putting the vial or prefilled diluent syringe in hot water.
- ▲ **Do not** put ALTUVIIIIO in direct sunlight.
- ▲ **Do not** return room temperature ALTUVIIIIO to the refrigerator.


Step 4:



Remove the plastic cap from the ALTUVIIIIO vial. Wipe the rubber stopper of the vial with an alcohol wipe and allow it to dry.

- ▲ After cleaning, **do not** touch the rubber stopper with your hand or allow it to touch any surface.


Step 5:



Completely remove the backing from the vial adapter package by peeling back the lid.

- ▲ **Do not** remove the vial adapter from the package or touch the inside of the vial adapter.

Step 6:




Keep the vial on a flat surface. Hold the vial with one hand and using the other hand, place the vial adapter in its package over the vial.

Note: The spike should be placed directly above the center of the rubber stopper.


- Push the vial adapter straight down until the spike on the vial adapter punctures the center of the vial stopper and is fully inserted.

Step 7:



Lift the package cover away from the vial adapter and throw away the cover.


Step 8:



Note: Only use the prefilled diluent syringe provided to mix (reconstitute) the powdered medicine.

- Hold the plunger rod by the circular disk.
- Place the tip of the plunger rod into the end of the prefilled diluent syringe.
- Turn the plunger rod to the right until it is firmly attached.

Step 9:




With one hand, hold the prefilled diluent syringe directly under the cap with the cap pointing up.

Note: Make sure you are holding the prefilled diluent syringe by the ridged part directly under the cap.

- ▲ **Do not** use if the cap has been removed or is not securely attached.

Step 10:




With your other hand, grasp the cap and bend it at a 90-degree angle until it snaps off.

Note: After the cap snaps off, you will see the glass tip of the prefilled diluent syringe.

- ▲ **Do not** touch the glass tip of the prefilled diluent syringe or the inside of the cap.


Step 11:



Note: Be sure the vial is sitting on a flat surface.

- Insert the tip of the prefilled diluent syringe into the vial adapter opening.
- Turn the prefilled diluent syringe to the right until it is securely attached to the vial adapter.

Step 12:



Slowly push down on the plunger rod to inject all of the liquid (diluent) from the prefilled diluent syringe into the vial.

Note: The plunger rod may rise slightly afterward. This is normal.

Step 13:

With the prefilled diluent syringe still connected to the adapter, gently swirl the vial until the powder is completely dissolved.


Check the solution through the vial to make sure the powder is fully dissolved.

Note: The solution should look clear and colorless to slightly yellowish.

▲ **Do not** shake.

▲ **Do not** use the mixed (reconstituted) ALTUVIIIIO if it contains visible particles or is cloudy.

POOLING: pooling is the process of combining two or more reconstituted vials into a larger luer lock syringe (not provided in the carton). If the dose requires more than one vial, reconstitute each vial as described above (See Steps 4–13) with the prefilled diluent syringe provided. **Do not** detach the prefilled diluent syringe until you are ready to attach the large luer lock syringe to the next vial. Keep the vial adapter attached to the vial as you will need it for attaching a large luer lock syringe. Use a larger luer lock plastic syringe to combine the contents of the reconstituted vials into the syringe, similar as described in the Steps 14–15. Repeat this pooling procedure with each vial you will be using. Once you have pooled the required dose, proceed with Step 16 (Injecting ALTUVIIIIO) using the larger luer lock syringe.

Step 14:

Make sure the plunger rod is pressed all the way down and the syringe is firmly attached to the vial adapter. Turn the vial upside-down.

Slowly pull down on the plunger rod to draw all the solution from the vial into the syringe.

Note: Be careful not to pull the plunger rod completely out of the syringe.

Step 15:

Gently unscrew the syringe from the vial adapter by turning it to the right. Throw away (dispose of) the vial with the adapter still attached (See Step 23). If you are not ready to inject, put the syringe cap carefully back onto the syringe tip.

▲ **Do not** touch the syringe tip or the inside of the cap.

Note: Mixed (reconstituted) ALTUVIIIIO should be given within **3 hours** after mixing.

Protect from direct sunlight. Do not refrigerate after reconstitution

Injecting ALTUVIIIIO

ALTUVIIIIO is given in the vein (intravenous injection) after mixing (reconstitution) of the powdered medicine with the diluent.

These are very generic instructions so your healthcare provider should teach you how to inject ALTUVIIIIO. After you have been taught to self-inject, you can follow these instructions.


▲ **Do not** give mixed (reconstituted) ALTUVIIIIO if it contains particles, is discolored, or is cloudy.

Step 16:


Using aseptic technique (clean and germ free), attach the syringe to the connector end of the infusion set tubing by turning it to the right until securely attached.

▲ **Do not** give mixed (reconstituted) ALTUVIIIIO in the same tubing or container with other medicines.

Note: Ask your healthcare provider which infusion set can be used with ALTUVIIIIO.


Step 17:

Apply a tourniquet to the upper arm. Then, using a new alcohol wipe, clean the skin where you will insert the needle and wait for it to dry.

Step 18:

Prime the syringe and the tubing. Push the plunger rod until all air is removed from the syringe and ALTUVIIIIO has filled the infusion set needle.


▲ **Do not** push ALTUVIIIIO through the needle.

Step 19:

Remove the protective needle cover from the infusion set needle and throw away (discard it).

▲ **Do not** touch the needle and do not put the needle cover back onto needle after use.

- Insert the needle on the infusion set tubing into the vein as instructed by your healthcare provider.

Step 20:

Use tape to secure the plastic wings of the needle in place at the injection site if needed.




Remove the tourniquet.

Note: Always make sure you have correctly inserted the needle into a vein when you perform an intravenous injection.

Initial approval: February 2023

AHF-FPLR-SL-FEB23

Rx Only

Step 21:	
	<p>Slowly push the plunger rod on the syringe all the way down to give ALTUVIIIIO.</p> <p>Your healthcare provider will provide your rate of administration based on your comfort level and the minimum injection time recommendation per vial.</p> <p>Note: A small amount of medicine will be left in the infusion set after injection. This is normal.</p>
Step 22:	
	<p>After delivering ALTUVIIIIO, remove the tape and the needle from the vein. Use a cotton ball or gauze pad to put pressure on the injection site for several minutes to stop any bleeding.</p> <p>Note: You may apply an adhesive bandage if needed.</p>
Step 23:	
	<p>Put the infusion set and the syringe in FDA-cleared sharps disposal container right away after use.</p> <p>Dispose of all unused solution according to your local regulations.</p> <p>Dispose of all empty vial(s), and other used medical supplies in your household trash.</p> <p>▲ Do not throw away (dispose of) the infusion set or syringe in your household trash.</p>
<p>Disposing of ALTUVIIIIO</p> <p>Throw away (dispose of) the infusion set and the syringe in FDA-cleared sharps disposal container right away after use.</p> <p>If you do not have FDA-cleared sharps disposal container you may use a household container that is:</p> <ul style="list-style-type: none"> - made of a heavy-duty plastic, - can be closed with a tight-fitting, puncture-resistant lid, without sharps being able to come out, - upright and stable during use, - leak-resistant, and - properly labeled to warn of hazardous waste inside the container. <p>When your FDA-cleared sharps disposal container is almost full, you will need to follow your community guidelines for the right way to dispose of FDA-cleared sharps disposal container. There may be state or local laws about how you should throw away used ALTUVIIIIO supplies.</p> <p>▲ Do not dispose of your used FDA-cleared sharps disposal container in your household trash unless your community guidelines permit this.</p> <p>▲ Do not recycle your used FDA-cleared sharps disposal container.</p> <p>Keep your FDA-cleared sharps disposal container out of the reach of children.</p>	

Manufactured by:
 Bioverativ Therapeutics Inc.
 Waltham, MA 02451
 A SANOFI COMPANY
 US License Number 2078